



CONTACT CENTER SCHEDULING USING INTEGER PROGRAMMING

5

Inventor: Turgut Aykin, Atlantic Highlands, NJ

Assignee: ac² Solutions, Inc.

1 Bethany Rd., Suite: 52

10 Hazlet, NJ 07730

taykin@ac2solutions.com

(732)768-3061

15

References Cited

U.S.PATENT DOCUMENTS

5,911,134	6/1999	Castonguay et al.,	364/401
20 6,044,355	3/2000	Crockett et al.	705/8
6,192,346	2/2001	Green	364/401
6,278,978	8/2001	Andre et al.	705/7

25

OTHER PUBLICATIONS

Danzig, G.B. , "A Comment on Edie's 'Traffic Delays at Toll Booths'," *Operations Research*, vol. 2, 339-341, 1954.

Holloran, T.J., and Byrn, J.E., "United Airlines Station Manpower Planning," *Interfaces*, vol. 16, 39-50, 1986.

Nanda, R., and Browne, J., *Introduction to Employee Scheduling*, Van Nostrand Reinhold, 1992.

5 Taha, H.A., *Operations Research*, 4th ed., Macmillan, Inc., 1987.

Wolsey, L.A., *Integer Programming*, John Wiley & Sons, 1998.

10

15

CONTACT CENTER SCHEDULING USING INTEGER PROGRAMMING

5

FIELD OF THE INVENTION

The invention pertains to the field of computer systems and computer
implemented methods. More specifically, the invention pertains to methods for
10 scheduling agents with multiple skills optimally at contact centers to satisfy time varying
agent staffing requirements for multiple contact types.

BACKGROUND OF THE INVENTION

5 The need for scheduling agents optimally in contact centers arises in a broad range of industry areas. Examples include call centers providing inbound or outbound services, telephone operator centers, catalog sales centers, hospitals, police stations, airports, maintenance and installation operations, manufacturing, etc. Agents may be call center operators, technicians, workers, servers, nurses, etc. Contact centers may include any service center providing services to its customers through their agents. Many
10 manufacturing operations may also fit into a contact center paradigm with similar needs and challenges in terms of worker or crew scheduling. Scheduling agents may be done a day at a time by specifying daily shift start time, shift length, time for the relief and lunch breaks, or to cover an entire week specifying, in addition to specifying daily shift schedules, work and non-work days. Contact centers use one or more interaction media
15 to provide the desired services to their customers. Interaction media (also referred to as “contact types”) may include telephone calls, web chat, email, fax, etc. It may also include contacts using different languages such as English, Spanish, etc., and/or different services such as sales, technical help, etc. Customers choose their time and media of contact to interact with the contact center agents. The contact type used by a customer
20 may require an agent with a special skill (e.g. language skills, writing proficiency). Moreover, agents may possess only some of the skills needed to serve customers using different contact types. Agents with the same skill set form a skill group (also referred to as “agent skill group”).

25 One of the critical tasks in contact center management is workforce scheduling. In such systems, workload changes during the course of an operating day, and from one day to another during a week. While understaffing will lower labor costs, it will result in poor service levels and longer waiting times for customers. Overstaffing, on the other hand, will improve service levels and lower waiting times, but will result in underutilized
30 agents and higher labor costs. Hence, it is important for contact center managers to schedule their workforce in an optimal manner to minimize labor costs while meeting the

projected agent and skills requirements. The present invention provides a method for formulating the workforce scheduling environment and requirements at a plurality of contact centers as a Mixed Integer Linear Programming (MILP) model, and a solution algorithm for developing optimal schedules for agents.

5

Traditionally, the task of developing work schedules to meet the forecasted agent and skill requirements is carried out manually. Manual scheduling when there is time varying workload and agent requirements, various daily shift and weekly tour scheduling rules, and multiple agent groups with different sets of skills and full-time/part-time work requirements is laborious and inefficient. Manpower scheduling in a typical contact center environment may involve an astronomical number of potential schedules, making manual evaluation of even a very small fraction of them to find an efficient schedule impractical.

15 In the past, a number attempts were made to develop more effective methods for workforce scheduling. Even though terms such as "optimal schedule", "optimized schedule" or "local optimal" were used in their descriptions, none of the prior art agent scheduling methods states or verifies the necessary and sufficient conditions for optimality in scheduling agent. Like the manual scheduling method, the prior art agent scheduling methods use a set of rules of thumb to develop agent schedules. None of them presents any evidence of, or guarantee the optimality of the schedules that may be obtained using the rules of their methods.

25 In the field of mathematical optimization, there are proven necessary and sufficient conditions for a solution to an optimization problem to satisfy in order to be the optimal (absolutely the best) solution (Taha, 1987, pages 743-779). The method of the present invention first develops an MILP model for which these necessary and sufficient conditions are well stated, and then uses an algorithm that would generate an optimal solution satisfying these conditions.

30

U.S. Pat. No. 5,911,134 issued on June 8, 1999 to Castonguay et al. describes a method for developing weekly tour schedules in call centers for agents with a single skill type. This method describes tour and shift construction rules to select tour and daily shift templates from the available ones, one day at a time, using measures such as “coverage”,
5 and break scheduling rules to schedule daily breaks (U.S. Pat. No. 5,911,134, col. 18, lines 26-46, and FIG. 11). The method also includes post-processing steps to eliminate the redundant tours, level break times, and take employee preferences into consideration. No evidence to suggest that the schedules obtained using these rules will satisfy the necessary and sufficient conditions for optimality in a workforce scheduling problem is
10 provided or known. As in the case of manual scheduling, the existence of an astronomical number of potential schedules, even in small call center scheduling environments, makes it very unlikely to locate the optimal schedule with the tour and shift construction rules used by this method.

U.S. Pat. No. 6,044,355 issued on March 28, 2000 to Crockett et al. describes a simulation method for developing weekly tour schedules in a contact center environment involving multiple agent groups with a plurality of skill sets, and a plurality of contact types requiring different agent skills. The method uses a scheduler and an Automatic Call Distributor (ACD) simulator. However, no working details or description of the
20 scheduler, and how it develops a schedule is included in the patent document. It was disclosed that the method uses a “scheduler program” (U.S. Pat. No. 6,044,355, col. 6, lines 18-36). No evidence to suggest that the schedules obtained using the scheduler program considered will satisfy the necessary and sufficient conditions for optimality in a workforce scheduling problem is provided or known.

U.S. Pat. No. 6,278,978 issued on August 21, 2001 to Andre et al. describes a method that may be used to post-process an initial schedule available from another source outside of the method by applying a rule-based interchange procedure. The method is provided for scheduling agents with a single skill type. The method begins by acquiring
30 an agent schedule developed by another method. The method then unschedules an agent from a shift in the schedule, evaluates the value of a “score” function for all other

schedules that may assign this agent to a different shift, and selects a shift using the score function to improve the schedule. The method also considers rescheduling of breaks, again using the score function. Interchanging two or more activities to improve the quality of an initial solution to a problem is a well-known strategy in the field of Optimization and the related research literature. It is also well known in the field of Optimization that interchanging (rescheduling) one or more agents and shifts simultaneously doesn't guarantee the optimality of the solution developed and the satisfaction of the optimality conditions.

A primary object of the present invention is to overcome these limitations of the prior art agent scheduling methods. The present invention uses Mixed Integer Linear Programming approach (MILP) (Wolsey, 1998) to develop a mathematical model of a workforce scheduling environment. A number of researchers including Danzig (1954) proposed MILP models for workforce scheduling. Difficulties with the use of the prior art MILP models for workforce scheduling are well documented (Nanda and Browne, 1992, page 206, Holloran and Byrn, 1986, page 13). The MILP model proposed by Dantzig (1954), for example, enumerates all possible combinations of shift and tour parameters resulting in tens of trillions of decision variables, making it very inefficient and impossible to solve even in small contact center environments. To overcome these limitations of the prior art MILP agent scheduling models, the present invention formulates daily break, and days-off scheduling implicitly (i.e. not explicitly enumerating all possible combinations). Thus, the method of the present invention formulates a significantly smaller but equally powerful MILP model, and uses various extensions of it.

A further limitation of the prior art MILP models is that they only consider agent scheduling environments involving a single agent skill group. Recent technological developments in ACD's used in call centers to queue and assign calls to agents, and incorporation of other types of contact media such as email and fax, made contact center managers to realize that agents with different types of skills are needed to be scheduled to handle these contact types (e.g. a Spanish speaking agent to answer a caller speaking in Spanish, or agents with good writing skills to respond to emails). Thus, the agent

scheduling task became more complex since the agent scheduling method used should take the time-varying demand for different agent skill types, and available agent groups with different skill sets (e.g. Spanish & English speaking sales, Chinese & English writing for technical help emails, and English only agent groups) into consideration in scheduling agents. It is another object of the present invention to disclose a method for formulating MILP models for scheduling environments involving a plurality of agent skill groups, and a plurality of contact types with time-varying workload for specific agent skills.

The present invention also discloses a solution algorithm that solves a number of sub-problems (or nodes) with the use of the standard Branch and Cut (B&C) algorithm for MILP problems. The solution method supplements the B&C algorithm by a Rounding Algorithm (RA algorithm) to locate the optimal solution of the MILP model of the invention. The optimality condition (both necessary and sufficient) for an MILP model is well documented in the field of Optimization; An all-integer solution that satisfies all of the constraints of an MILP with a minimization (maximization) type objective function, and has an objective value that is better than the best lower bound (upper bound) for any of the sub-problems in the B&C algorithm is a global (i.e. absolute) optimal solution to that MILP model (Wolsey, 1998). Once the optimality condition is satisfied by an all-integer solution found during the execution of this solution algorithm (either in the B&C algorithm or in the RA algorithm), an optimal agent schedule is reported using the information in the optimal solution of the MILP model found.

It is still another object of the present invention to provide a computer implemented optimal workforce scheduling method for contact centers that schedule their workforce to meet varying workload during the course of a day, and from one day to another during a week.

SUMMARY OF THE INVENTION

The method of the present invention first formulates a skills based or non-skills based agent scheduling environment involving a plurality of agent skill and work groups, and plurality of contact types as an MILP model including various contact center constraints and requirements. The formulation step of the present invention involves, first, the definitions of various decision variables, and parameters, and the formulation of an objective function. The objective function of the MILP model of the present invention determines a measure of effectiveness (or merit) for each solution. Examples of the objective function in the method include total schedule cost, total time scheduled, total paid time scheduled, total agent preference, etc. The formulation step also includes the expression of contact center constraints, and agent and skill requirements in all periods to be scheduled as mathematical equalities or inequalities. Finally, constraints that restrict the values of all decision variables to nonnegative values, and some variables to nonnegative integer values only are added to the formulation.

The MILP model of the present invention may have one or more solutions (also called "feasible solutions") that satisfy all contact center constraints and requirements included in it. The optimal solution of an MILP model is merely finding the best feasible solution in terms of its objective function value among *all* feasible solutions. The best feasible solution with the most favorable objective function value is the optimal solution and satisfies the optimality condition. Moreover, every agent schedule satisfying the optimality condition in a specific workforce environment, when all contact center constraints and requirements in each planning period are satisfied, is also a global or absolute optimal solution (e.g. see Taha, 1987). The optimal agent schedule is then reported using the optimal solution.

To find a global optimal solution to the MILP model of an agent scheduling environment, the present invention applies an enhanced optimization algorithm combining the standard technique for solving MILP models optimally, the B&C algorithm (Wolsey, 1998), with an RA algorithm. An MILP model contains one or more

decision variables whose values are restricted to non-negative integer values. In the case of the MILP of the present invention, for example, the number of agents assigned to a weekly work pattern specifying their work and non-work days can't take a fractional value (e.g. it can't be 4.75 agents). The B&C algorithm first solves the MILP model (called "node zero") without integrality constraints for these decision variables. This version of the MILP model is referred to as the Linear Programming (LP) relaxation of the MILP. If the values of the decision variables in this solution also satisfy the integrality constraints for all decision variables, the solution is optimal. The solution algorithm of the present invention then terminates, and the schedule-reporting step begins.

If the values of the decision variables in the solution of the LP relaxation at node zero don't satisfy all integrality constraints, then the solution algorithm of the present invention calls the RA algorithm to search for an integer feasible solution. The RA algorithm first retrieves the values of the decision variables found by the B&C algorithm to the LP relaxation. The RA algorithm then rounds the values of the integer restricted decision variables with fractional values either up or down in an attempt to find an integer feasible solution. After rounding the values of the decision variables, the RA algorithm also checks the constraints of the MILP model. If the rounded solution is violating one or more constraints of the MILP model, values of the variables are adjusted to restore feasibility including addition of new agents to meet the agent requirements in every period, and for all skill types. If no solution satisfying all constraints can be found with the current solution, the RA algorithm terminates and control is returned back to the B&C algorithm. Otherwise, if an integer feasible solution to the MILP model is found, the RA algorithm calculates the objective value for that solution, and compares it with the best known solution to the MILP model, if one is available. If the new integer feasible solution found has a more favorable objective value, the new integer feasible solution replaces the best known solution to the MILP. The RA algorithm then terminates, and returns the new integer solution found back to the B&C algorithm.

30

If the best known (integer feasible) solution was changed in the RA algorithm, the B&C algorithm updates the best integer solution known for its own use, and forms new sub-problems (nodes) by adding new constraints (Wolsey, 1998). The entire process is then repeated; the B&C algorithm chooses a new sub-problem, and solves its LP

5 relaxation. If the LP relaxation has a feasible solution satisfying all of the constraints but some of the integrality constraints for some decision variables, the B&C algorithm transfers control to the RA algorithm which searches for an integer feasible solution through rounding and adding more agents to the schedules. The solution algorithm of the present invention terminates when all sub-problems in the B&C algorithm are terminated
10 (sub-problem termination conditions in the B&C algorithm are given in Wolsey, 1998). That is when the optimality condition is satisfied by an integer feasible solution to the MILP model. When this condition is satisfied, control is transferred to the schedule-reporting step.

15 In the schedule reporting step of the present invention, an optimal agent schedule with daily start and stop times, work and non-work days, daily break times for every work day, agent allocation to contact types in every period to be scheduled are reported for every agent tour schedule.

20

BRIEF DESCRIPTION OF THE DRAWINGS

For a more complete description of the present invention and its merits, reference should be made to the Description of the Preferred Embodiment.

5

FIG. 1 shows the process architecture for optimal workforce scheduling with the method of the present invention, and its computer implementation.

FIG. 2 shows a flowchart of the MILP model generation of the invention.

FIG. 3 shows all possible work patterns in a week.

10

FIG. 4 shows a flowchart of the solution method for the MILP model of the invention.

DESCRIPTION OF THE PREFERRED EMBODIMENT

FIG. 1 shows process architecture for optimal workforce scheduling with the method of present invention, and its computer application. It consists of several sub-processes including forecasting (1), agent and skills requirements planning (2), scheduling environment information input (3) (e.g. center information, agent skills and work group availability, tour and shift parameters), MILP model generator (4), optimizer (5), and rostering (6) processes. The main function of the forecast generator is to make predictions of future contact volumes, and average handling times (service times) for the contact types offered in each planning period during the days in the schedule interval (e.g. a week). This sub-process uses statistical techniques to analyze patterns in the historical data, and develop forecasts. Agent and skill requirements planning module (2) combines contact volume and average handling time forecasts with service level targets (e.g. 80% of incoming calls answered within 20 seconds of their receipt) to determine agent and skill requirements in each planning period to be scheduled. In a multi-skilled environment, this module determines the agent and skills mix from the available agent skill groups that are needed in each planning period to deliver the targeted service levels to all contact types. This process also takes the economies of multi-skilled scheduling environment into consideration. Contact center managers input the environment information including the objective criterion to be optimized, and various contact center parameters (e.g. number, location, operating hours), agent skill and work groups, and tour and shift parameters to module (3). The next two modules, the MILP model generator (4) and optimizer (5), use the method of present invention to develop optimal agent schedules to meet agent and skills requirements developed in module (2) using the scheduling environment parameters provided in module (3). The present invention provides a method for the MILP model generation for a workforce scheduling environment, and an algorithm for a near-optimal or optimal solution of the MILP model generated by the MILP model generator to develop agent schedules. The method of the present invention covers both non-skills based, and skills based agent scheduling environments.

The scheduling environment may involve one or more contact centers. In the case of multiple contact centers, each contact center may be scheduled independently to meet their allocated agent and skills requirements with their own agent skill groups, agent work groups, agent availability, operating hours, and tour and shift scheduling rules. In this case, the method of the invention formulates the scheduling environment at each contact center as an MILP model and solves it optimally to obtain an optimal schedule. Alternatively, a plurality of contact centers may be scheduled as a virtual contact center. In this case, agent and skills requirements of different contact centers are combined using a common time zone, and the method of invention formulates one MILP model for these centers to include combined agent and skills requirements with the centers' agent skill groups, agent work groups, agent availability, operating hours, and tour and shift scheduling rules. When solved together, the optimal schedule provides a combined schedule for agents at all of the centers included. The unique operating characteristics of each contact center such as operating hours, tours and shift scheduling rules, agent skill and work group availability, and agent availability are satisfied by modeling them in one common time zone, and identifying them by contact center specific identification numbers.

FIG. 2 shows the basic steps of the MILP model generation method of the present invention. The following description, referring to FIG. 2, is presented to explain the MILP model generation method in its general case, before proceeding to the more specific scheduling environments.

Step 7: Acquire agent and skills requirements for each planning period to be scheduled.

Step 8: For each contact center, obtain daily operating hours, agent skill groups, work groups for these agent skill groups specifying various work patterns such as full-time 40 hours/week, or part-time 30 hours/week, and the scheduling paradigm as to whether to schedule each contact center separately or a plurality of them in a virtual contact center environment.

Step 9: Acquire tour and shift parameters and scheduling rules, and their availability to

agent skill and work groups at different contact centers.

Step 10: Acquire the objective criterion to be optimized and associated cost and benefit parameters.

Step 11: Generate the appropriate MILP model by first defining decision variables, parameters and sets, and then formulating the objective function, agent and skill requirement constraints, break constraints, days-off and work pattern constraints, agent availability constraints, other schedule related constraints, and non-negativity and integrality constraints.

Step 12: Input the variables, objective function, and constraints of the MILP model formulated to optimizer (2).

A number of MILP formulations of the present invention for agent scheduling in a non-skills based (i.e. single skill) environment and their extensions are discussed first. The MILP model generation method of the present invention for skills based agent scheduling environments involving agents with different skills serving multiple contact types is disclosed after this discussion. The method of the invention for optimal solution of the MILP model is disclosed thereafter.

To facilitate the description of the MILP model and its extensions used by the present invention in this patent document, any variable or set defined with a subscript is shown as flat text when the variable or set itself is a subscript of another term. For instance, a set defined as B_{1kih} is typed as B_{1kih} in $\sum_{t \in B_{1kih}}$ where it is a subscript of the summation sign \sum . Moreover, symbols “ ε ” and “ \in ” are both used to imply membership in a set or indicate an element of a set.

**Workforce Scheduling with Tours Requiring Consistent
Daily Start Time and Shift Length in
a Single Skill Environment**

5 To facilitate the presentation, this discussion assumes that all agents have the same skill set (i.e. not a skills based scheduling environment). In addition, the following assumptions regarding the scheduling environment are made to facilitate the presentation: Each agent works 5 days and receives two consecutive days-off during a week; There are one or more contact centers operating 24 hours or less a day and 7 days a week. A day is
10 divided into equal planning periods (e.g. 15-minute intervals) that are indexed sequentially in T_h for day h , $h = 1, \dots, 7$. As disclosed below, none of these specifications is restrictive on the MILP model of the method. A person with ordinary skills at art can easily extend the MILP model of the method to cases where the number of work days may be less or greater than 5 days per week, a work tour less than or
15 exceeding 7 days, any number of consecutive days-off, and a scheduling environment where the time period may be less or greater than 15 minutes.

Moreover, for this discussion, it is also assumed that a tour specifies a weekly work pattern that is seven days long with the following breaks in the work schedule: (i) Daily
20 breaks in the shift schedule including two relief breaks, and one lunch break during a work day, and (ii) two consecutive non-work days (i.e. days-off). We assume that each relief break is one planning period, and lunch break is two planning periods long. Extensions of the MILP model of the method to tours with more or less breaks, and different break durations are discussed with other cases below. Agents assigned to a tour
25 have the identical daily shift start times and shift lengths during a week. The present invention extends the MILP model to the scheduling environments involving non-identical daily shift start times and shift lengths later in this document. The scheduling environment may include more than one tour type. Each tour type specifies a set of daily shift start times, and a shift length for seven days. In the general case, different tour types
30 may also have different number of consecutive or non-consecutive off days, number of daily breaks and break durations.

Let K be the set of all distinct tour types available to assign agents. Each tour type k has a predetermined set of start times given in I_k . Let a_{kht} be equal to one if period t on day h is in the shift span (that is, a work, break, or days-off period) of tour k with a daily start time of i , and zero otherwise. Suppose that the maximum number of agents available to assign to tour k is limited to D_k^{\max} . Let X_{ki} be the number of agents assigned to tour k with a daily start time of period i . Let C_{ki} be the weekly cost (wages and benefits, etc.) of assigning an agent to tour k with a daily start time of i , $i \in I_k$, and $k \in K$. Let U_{kiht} , W_{kiht} and V_{kiht} , daily break variables, be the numbers of agents assigned to tour k with a daily start time of i and starting their first relief, lunch and second relief breaks in period t on day h , respectively. Daily relief and lunch breaks must start and end within the break windows specified for them. Let $B1_{kih}$, $B2_{kih}$ and $B3_{kih}$ be the sets of planning periods on day h during which an employee assigned to tour k with a daily start time of i may start, respectively, his/her first relief, lunch and second relief break and complete within the time windows specified for these breaks. Break variables $U_{kih(t1)}$, $W_{kih(t2)}$ and $V_{kih(t3)}$ are defined for $t1 \in B1_{kih}$, $t2 \in B2_{kih}$, $t3 \in B3_{kih}$, and $i \in I_k$, $k \in K$, $h = 1, \dots, 7$, respectively. Further, let $T1_{kht}$, $T2_{kht}$ and $T3_{kht}$ be the sets of start times for tour k for which period t on day h is, respectively, a first relief, lunch, and second relief start period within the break windows specified for them. Let Y_{kih} be the number of agents assigned to tour k with a daily start time of i who are scheduled start their consecutive non-work days (days-off) on day h . Finally, let b_{ht} be the number of agents required in period t on day h to deliver the desired service level to customers. The MILP model of the present invention for the scheduling environment assumed can be stated as follows.

$$\text{Minimize } \sum_{k \in K} \sum_{i \in I_k} C_{ki} X_{ki} \quad (1)$$

Subject to

$$\begin{aligned} 5 \quad & \sum_{k \in K} \sum_{i \in I_k} a_{kiht} X_{ki} - \sum_{k \in K} \sum_{i \in I_k} a_{kiht} (Y_{kih} + Y_{ki(h-1)}) \\ & - \sum_{k \in K} \sum_{i \in T1kht} U_{kiht} - \sum_{k \in K} \sum_{i \in T2kht} (W_{kiht} + W_{kih(t-1)}) \end{aligned}$$

$$- \sum_{k \in K} \sum_{i \in T3kht} V_{kiht} \geq b_{ht},$$

10

$$t \in T_h, h = 1, \dots, 7, \quad (2)$$

$$X_{ki} - Y_{kih} - Y_{ki(h-1)} = \sum_{t \in B1kih} U_{kiht}, \quad i \in I_k, k \in K, h = 1, \dots, 7, \quad (3)$$

15

$$X_{ki} - Y_{kih} - Y_{ki(h-1)} = \sum_{t \in B2kih} W_{kiht}, \quad i \in I_k, k \in K, h = 1, \dots, 7, \quad (4)$$

$$X_{ki} - Y_{kih} - Y_{ki(h-1)} = \sum_{t \in B3kih} V_{kiht}, \quad i \in I_k, k \in K, h = 1, \dots, 7, \quad (5)$$

20

$$X_{ki} = \sum_h Y_{kih}, \quad i \in I_k, k \in K, \quad (6)$$

$$\sum_{i \in I_k} X_{ki} \leq D_k^{\max}, \quad k \in K, \quad (7)$$

25

$$X_{ki}, Y_{kih}, U_{kiht}, W_{kiht}, \text{ and } V_{kiht} \geq 0, \text{ and integer for all } k, i, h, \text{ and } t. \quad (8)$$

Objective function (1) is the total cost of assigning agents to various tour types and start times per week. Constraint (2) ensures that the total number of agents assigned to various tours and available in period t on day h (first term on the left side of (2)) minus the number of agents who are assigned to these tours but taking this day off (second term on the left side of (2)) and the number of agents who are at work on day h but taking their first relief (third term on the left side of (2)), lunch (fourth term on the left side of (2)), and second relief (fifth term on the left side of (2)) in period t on day h is sufficient to meet the agent requirement, b_{ht} , in that period. The second term on the left side of (2) includes agents who are starting their consecutive days-off schedule on day h as well as those who started a day earlier on day $(h-1)$. Break constraints (3-5) ensure that one first relief, one lunch, and one second relief break are scheduled for each agent who is assigned to tour k with a daily start time of i and working on day h . Thus, the number of first relief, lunch, and second relief breaks scheduled for tour k with a daily start time of i on a given day is equal to the number of agents assigned to this tour and start time minus the number of them scheduled to take day h off, $X_{ki} - Y_{kih} - Y_{ki(h-1)}$ (because of two consecutive days-off per week requirement). Constraint (6) ensures that a set of two consecutive days-off is scheduled for every agent assigned to tour k with a daily start time of i . Constraint (7) ensures that the total number of agents assigned to tour type k doesn't exceed the maximum number of agents that are available to assign to this tour type. This constraint can easily be modified to limit the number of agents that can be assigned to a set of tour types (e.g. full time 40-hour a week tours may include 8-hour shifts and 5 work days a week, and 10-hour shifts and 4 work days a week). Finally, constraint (8) restricts the tour variables X_{ki} , days-off variables Y_{kih} , and break variables U_{kiht} , W_{kiht} , and V_{kiht} to nonnegative integers to avoid negative values and/or fractional agent schedules.

Extensions of MILP Model (1-8)

It should be clear to those skilled in the art that the specific embodiments and the MILP model disclosed above may be readily utilized as a basis and extended to other

scheduling environments without departing from the scope of the invention. For example, objective function (1) can be easily reformulated to minimize the total amount of agent time scheduled or paid agent time scheduled by setting C_{ki} equal to scheduled or paid time per week per agent assigned to tour k with a daily start time of i , $i \in I_k$, and $k \in K$.

5 Also, if agents rate their most preferred, second most preferred, etc., tours using an RAting system, these ratings for various tours and start times can be aggregated and used in the objective function to maximize agent preference in scheduling tours and start times. Note that in this case, in order to not to schedule redundant tours (a redundant tour is a tour that can be removed without causing agent shortages in any of the periods
10 covered by that tour during the week) and increase agent costs unnecessarily, rating system should assign the lowest value to the most preferred tour, a higher to the second most preferred tour, etc. Consequently, the objective function will remain as of minimization type to maximize agent preferences.

15 Another extension of MILP model (1-8) involves agent scheduling when there aren't enough agents to meet the agent requirements in every period to be scheduled. In this case, objective function (1), and constraint (2) are modified by adding overstaffing and understaffing variables. Let O_{ht} be the number of agents in excess of agent requirements, and S_{ht} be the agent shortage in period t on day h . A penalty for shortages
20 in a period, P_{ht} , is included in the objective function to find the optimal schedule that minimizes the total agent cost plus the shortage cost.

$$\text{Minimize } \sum_{k \in K} \sum_{i \in I_k} C_{ki} X_{ki} + \sum_h \sum_{t \in T_h} P_{ht} S_{ht} \quad (9)$$

25 Constraint (2) is then formulated by adding O_{ht} for overstaffing and S_{ht} for shortages to its left side as follows:

$$\begin{aligned} & \sum_{k \in K} \sum_{i \in I_k} a_{kiht} X_{ki} - \sum_{k \in K} \sum_{i \in I_k} a_{kiht} (Y_{kih} + Y_{ki(h-1)}) \\ & - \sum_{k \in K} \sum_{i \in T1kht} U_{kiht} - \sum_{k \in K} \sum_{i \in T2kht} (W_{kiht} + W_{kih(t-1)}) \end{aligned}$$

$$- \sum_{k \in K} \sum_{i \in T3kht} V_{kiht} + S_{ht} - O_{ht} = b_{ht},$$

$$t \in T_h, h = 1, \dots, 7,$$

5

(10)

$$S_{ht}, O_{ht} \geq 0, \text{ for all } t \in T_h \text{ and } h.$$

10 If a certain amount of overstaffing is needed for other tasks (e.g. to answer email or fax based contacts), it can be ensured by restricting the overstaffing to be greater than or equal to the desired level f_{ht} .

$$\sum_{t \in T_h} O_{ht} \geq f_{ht}, \quad t \in T_h, h = 1, \dots, 7, \quad (11a)$$

15

If up to a certain amount of understaffing staffing is tolerable (e.g. to see if the total cost can be reduced), this can be achieved by adding a similar constraint:

$$\sum_{t \in T_h} S_{ht} \geq d_{ht}, \quad t \in T_h, h = 1, \dots, 7, \quad (11b)$$

20

where d_{ht} is the maximum understaffing tolerated on day h . Constraints (11a) and (11b) can easily be further modified to impose limits for a specific period(s) on day h .

25 If a tour type has more than three breaks on a given day (e.g. first relief, lunch, second, and third relief breaks on Thursdays and Fridays), the method of the present invention extends MILP model (1-8) by introducing a set of break variables and a break constraint (similar to constraints (5-7)) for every break that needs to be scheduled so that every agent assigned to tour k and working on that day will have the correct number of breaks scheduled for him/her. If agents assigned to a tour are given fewer breaks (e.g. part time agents working 4 hours a day), the method of the present invention extends

MILP model (1-8) by excluding the break variables and constraints associated with the breaks not allowed.

Furthermore, if a tour type has a break longer than 2 consecutive planning
 5 periods, then the method of the present invention extends MILP model (1-8) by
 subtracting all agents who started their breaks in an earlier period but haven't completed
 yet and the agents who are starting their break in period t from the left side of constraint
 (2). For a 4-period break, for example, agents who started their breaks three periods
 earlier, two periods earlier, one period earlier, and agents starting in the current period
 10 will be subtracted from the number of agents scheduled in that period from the left side of
 constraint (2).

In MILP model (1-8), even though daily shift start times for an agent remain the
 same on different days, break times may vary from one work day to another. For
 15 instance, it is possible to have the first relief break of an agent be scheduled at 9:00am on
 Monday, and at 10:00am on Tuesday. If a tour (say tour k) requires consistent relief and
 lunch break times on all work days in addition to consistent daily shift start times and
 shift lengths, these requirements are ensured by adding the following constraints the
 MILP model (1-8):

20

$$\sum_h U_{kiht} \leq q_k SU_{kit}, \quad , i \in I_k, t \in D1_{ki}, \quad (12a)$$

$$\sum_h W_{kiht} \leq q_k SW_{kit}, \quad , i \in I_k, t \in D2_{ki}, \quad (12b)$$

$$\sum_h V_{kiht} \leq q_k SV_{kit}, \quad , i \in I_k, t \in D3_{ki}, \quad (12c)$$

$$25 \quad \sum_{t \in D1_{ki}} SU_{kit} = 1, \quad , i \in I_k, k \in K, \quad (13a)$$

$$\sum_{t \in D2_{ki}} SW_{kit} = 1, \quad , i \in I_k, k \in K, \quad (13b)$$

$$\sum_{t \in D3_{ki}} SV_{kit} = 1, \quad , i \in I_k, k \in K, \quad (13c)$$

$SU_{kit}, SW_{kit}, SV_{kit} \geq 0$ and integer.

where $D1_{ki}$, $D2_{ki}$, and $D3_{ki}$ are the sets containing the first consistent (across the days of a week) daily first relief, lunch, and second break times in $B1_{kih}$, $B2_{kih}$, and $B3_{kih}$, respectively, q_k is a very large number, SU_{kit} , SW_{kit} , and SV_{kit} are binary variables taking
 5 a value of either zero or one to indicate whether period t is selected as the consistent daily first relief, lunch, or second relief break time, respectively.

Constraints (12a) and (13a) are restricting the daily first relief break times to one of the break times specified in $B1_{kih}$. Constraint (13a) ensures that only one relief break
 10 time is selected, that is, only one SU_{kit} is equal to one while the others are zero. Thus, for example, if the first relief break times for a tour starting at 7:00am daily are $B1_{kih} = \{8:15\text{am}, 8:30\text{am}, 8:45\text{am}\}$, $h = 1, \dots, 7$, then only one of $SU_{ki(8:15)}$, $SU_{ki(8:30)}$, or $SU_{ki(8:45)}$ will be equal to one and the others will be equal to zero. Likewise, constraints (12b) and (13b) are restricting the lunch break times, and constraints (12c) and (13c) are restricting
 15 the second break times to consistent daily schedules.

MILP model (1-8) of the method is presented for tour types that require two consecutive non-work days be scheduled for each agent. The present invention extends MILP model (1-8) to include tours with 3 or more consecutive days-off by subtracting
 20 from the left sides of constraints (2-5) the agents assigned to this type of tours and who are off as part of their scheduled consecutive non-work days, and agents starting their days-off on day h .

If, however, tours in K require a minimum of two consecutive days off and, in
 25 addition, a total of r_k non-consecutive days off, the present invention extends MILP model (1-8) as follows:

$$\text{Minimize } \sum_{k \in K} \sum_{i \in I_k} C_{ki} X_{ki} \quad (14)$$

Subject to

$$5. \quad \sum_{k \in K} \sum_{i \in I_k} a_{kiht} X_{ki} - \sum_{k \in K} \sum_{i \in I_k} a_{kiht} (Y_{kih} + Y_{ki(h-1)})$$

$$- \sum_{k \in K} \sum_{i \in I_k} \sum_{m \neq h, (h-1)} a_{kiht} Z_{kimh} - \sum_{k \in K} \sum_{i \in T1kht} U_{kiht}$$

$$- \sum_{k \in K} \sum_{i \in T2kht} (W_{kiht} + W_{kih(t-1)}) - \sum_{k \in K} \sum_{i \in T3kht} V_{kiht} \geq b_{ht}$$

10

$$, t \in T_h, h = 1, \dots, 7, \quad (15)$$

$$X_{ki} - Y_{kih} - Y_{ki(h-1)} - \sum_{m \neq h, (h-1)} Z_{kimh} = \sum_{t \in B1kih} U_{kiht}, \quad i \in I_k, k \in K, h = 1, \dots, 7, \quad (16)$$

$$15 \quad X_{ki} - Y_{kih} - Y_{ki(h-1)} - \sum_{m \neq h, (h-1)} Z_{kimh} = \sum_{t \in B2kih} W_{kiht}, \quad i \in I_k, k \in K, h = 1, \dots, 7, \quad (17)$$

$$X_{ki} - Y_{kih} - Y_{ki(h-1)} - \sum_{m \neq h, (h-1)} Z_{kimh} = \sum_{t \in B3kih} V_{kiht}, \quad i \in I_k, k \in K, h = 1, \dots, 7, \quad (18)$$

$$r_k Y_{kih} = \sum_{l \neq h, (h+1)} Z_{kihl}, \quad i \in I_k, k \in K, h = 1, \dots, 7, \quad (19)$$

20

$$Z_{kihm} \leq Y_{kih}, \quad i \in I_k, k \in K, h = 1, \dots, 7, \\ m \neq h, (h+1) \quad (20)$$

$$X_{ki} = \sum_h Y_{kih}, \quad i \in I_k, k \in K, \quad (21)$$

$$25 \quad \sum_{i \in I_k} X_{ki} \leq D_k^{\max}, \quad k \in K, \quad (22)$$

$$X_{ki}, Y_{kih}, Z_{kihm}, U_{kiht}, W_{kiht}, \text{ and } V_{kiht} \geq 0 \text{ and integer for all } k, i, h, m, \text{ and } t. \quad (23)$$

where Z_{kihm} is defined as the number of agents who are assigned to tour k with a daily start time of i and will start their consecutive days off on day h and will also take day m off as part of their tour schedule. Note that Z_{kihm} is defined for $m \neq h, (h+1)$, for tours requiring a minimum of two consecutive days off, in order to avoid infeasible days off schedules.

Constraint (19) ensures that each agent assigned to tour k receives exactly r_k additional days off. If for tour k , a maximum of $(2 + r_k)$ days-off with a minimum of 2 consecutive days-off, this can be formulated by simply modifying constraint (19) as a greater than or equal to type constraint (that is, the term on the left side of (19) is greater than or equal to the term on the right side).

Constraints (19) and (20) ensure that r_k different days are scheduled as non-work days for every agent on days other than their consecutive days off. In constraints (15-18), agents who are off on day h due to their consecutive days-off schedule as well as the agents who are off due to their additional days off schedule are subtracted from the total number of agents scheduled.

When a scheduling environment involves only tour types not requiring any consecutive days off, that is equivalent to requiring a minimum of one "consecutive" day off. In this case, the present invention extends MILP model by removing Z_{kihl} variables from all constraints, and dropping constraints (19) and (20). Y_{kih} is now defined as the number of agents assigned to tour k with a daily start time of i who will take day h off. Note that only Y_{kih} will be in constraints (15-18) instead of both Y_{kih} and $Y_{ki(h-1)}$ since the tour types considered do not require consecutive days off. The MILP model is now stated as follows:

$$\text{Minimize } \sum_{k \in K} \sum_{i \in I_k} C_{ki} X_{ki} \quad (24)$$

Subject to

$$5 \quad \sum_{k \in K} \sum_{i \in I_k} a_{kiht} X_{ki} - \sum_{k \in K} \sum_{i \in I_k} a_{kiht} Y_{kih} - \sum_{k \in K} \sum_{i \in T1kht} U_{kiht}$$

$$- \sum_{k \in K} \sum_{i \in T2kht} (W_{kiht} + W_{kih(t-1)}) - \sum_{k \in K} \sum_{i \in T3kht} V_{kiht} \geq b_{ht}$$

$$, t \in T_h, h = 1, \dots, 7, \quad (25)$$

10

$$X_{ki} - Y_{kih} = \sum_{t \in B1kih} U_{kiht} \quad , i \in I_k, k \in K, h = 1, \dots, 7, \quad (26)$$

$$X_{ki} - Y_{kih} = \sum_{t \in B2kih} W_{kiht} \quad , i \in I_k, k \in K, h = 1, \dots, 7, \quad (27)$$

15

$$X_{ki} - Y_{kih} = \sum_{t \in B3kih} V_{kiht} \quad , i \in I_k, k \in K, h = 1, \dots, 7, \quad (28)$$

$$r_k X_{ki} = \sum_h Y_{kih} \quad , i \in I_k, k \in K, \quad (29)$$

$$\sum_{i \in I_k} X_{ki} \leq D_k^{\max} \quad , k \in K, \quad (30)$$

20

$$X_{ki}, Y_{kih}, U_{kiht}, W_{kiht}, \text{ and } V_{kiht} \geq 0 \text{ and integer for all } k, i, h, \text{ and } t. \quad (31)$$

where r_k is now defined as the number of days-off an agent assigned to tour k receives.

25

A tour is said to have a fixed day off if every agent assigned that tour will be off on the same day. For example, a contact center may close on Sundays. In this case, all agents will receive a day off on Sunday. The present invention extends the MILP model

by setting the days-off variables for the fixed days off equal to the number of agent assigned to that tour. For example, if tour k requires two consecutive days off, and day h is an off day for all agents assigned to this tour, then

$$X_{ki} = Y_{kih} + Y_{ki(h-1)}, \quad i \in I_k, \quad (32)$$

And if tour k doesn't require consecutive days off, and day h is an off day for all agents assigned to this tour, then

$$X_{ki} = Y_{kih}, \quad i \in I_k. \quad (33)$$

If operating hours are shorter for certain days (e.g. regular hours of 5:00am to 10:00pm with Saturday operating hours of 5:00am to 4:00pm) for a tour or a contact center, the present invention extends the MILP model by formulating each such tour as consisting of two tours with identical work rules except for their start times as follows: (i) tour 1 includes the start times for which the latest daily shift completion time not exceeding the operating hours for the early closure day, and (ii) tour 2 has the start times not covered by tour group 1 (ending after closure on the early closure day) and has a fixed day off on the early closure day to not to schedule any agent to work on those days.

A scheduling environment can have any combination of the extensions and tour types discussed above. The MILP model of the present invention can be extended to formulate these scheduling environments by combining various extensions of the constraints and objective function together with the definitions of decision variables and parameters disclosed above.

**Workforce Scheduling with Tours not Requiring Consistent
Daily Start Time and Shift Length in
a Single Skill Environment**

5

To facilitate the discussion and notation, the scheduling environment described before the disclosure of MILP model (1-8) is considered again. The present invention also extends the MILP model to tours not requiring consistent daily shift start times and same daily shift length requirements. For these tours, shift start time as well as shift
10 duration for an agent may vary from one day to another.

Assume that the scheduling environment involves only tours not requiring consistent daily start time and shift length. Let QK be the set of all such tour types. Assume that agents assigned to tour type k can start at any one of a set of predetermined
15 start times given in QI_k . Let the daily shift lengths allowed for tour type k be F_k . F_k may also change from one day to another. Note that tour k has a minimum weekly work limit specified by the shortest shift in F_k , and a maximum weekly work limit specified by the longest shift in F_k . The daily shift start times should not allow for back-to-back shift
20 schedules for agents without allowing enough time for rest. Thus, the latest start time for a tour plus the length of the longest daily shift type are not allowed to exceed 24 hours minus the desired minimum rest period between consecutive daily shifts. To prevent this for a tour type, the present invention creates separate tour types each with start times allowing a minimum required time between the latest possible end of a shift on one day and the earliest possible start of a shift on the following day.

25

Moreover, assume that the maximum number of agents available to assign to tour k is QD_k^{\max} . Let QX_{knh} be the number of agents assigned to tour k and shift length n with a starting of period i on day h . Define daily break variables QU_{knh} , QW_{knh} and QV_{knh} as the numbers of agents assigned to tour k and shift length n with a daily start
30 time of i and starting their first relief, lunch and second relief breaks in period t on day h , respectively. Depending on the shift length, some of the breaks may not be provided.

This case is formulated in the MILP model as disclosed before for tours with more/fewer daily breaks. Daily relief and lunch breaks must start and end within the break windows specified for them. Let $QB1_{knh}$, $QB2_{knh}$ and $QB3_{knh}$ be the sets of planning periods on day h during which an employee assigned to tour k and shift length n with a daily start time of i may start his/her first relief, lunch and second relief break and complete within the time windows specified for these breaks, respectively. Break variables $QU_{knh(t1)}$, $QW_{knh(t2)}$ and $QV_{knh(t3)}$ are defined for $t1 \in QB1_{knh}$, $t2 \in QB2_{knh}$, and $t3 \in QB3_{knh}$, respectively. Further, let $QT1_{knh}$, $QT2_{knh}$ and $QT3_{knh}$ be the sets of start times for shift length n for tour k for which period t on day h is a first relief, lunch, and second relief start period, respectively. Let a_{knhit} be equal to one if period t on day h is in the shift span (that is, a work or a break period) of agents assigned to tour k who have the start time of i and shift length n on day h , and zero otherwise. A work pattern in this case is a string of zeros and ones indicating work days ($=1$) and non-work days ($=0$) scheduled for agents assigned to this work pattern. All possible work patterns for a week are shown in FIG. 3

Let the set of all allowed work and non-work patterns for the agents assigned to tour k be QL_k . Only the work patterns in FIG. 3 satisfying the work and non-work day rules specified for tour k are included in QL_k (e.g. a minimum of two consecutive days off, no work on Sunday, etc.). A work pattern in this case specifies work and non-work days explicitly. Define the days-off pattern variable Q_{kl} as the number of agents assigned to tour k with a work pattern of l . Let A_{klh} be equal to one if day h is a work day for agents assigned to work pattern l of tour k , and zero otherwise.

Let C_{kl} be the weekly "fixed" cost (e.g. benefits, etc.) of assigning an agent to tour k with work pattern of l , $l \in QL_k$, and $k \in QK$. Let c_{kni} be the daily wage paid to agents assigned to tour type k with a daily shift length of n and start time of i . Finally, let b_{ht} be the number of agents required in period t on day h to deliver the desired service level to customers. The present invention extends the MILP model to this scheduling environment as follows.

$$\text{Minimize } \sum_{k \in QK} \sum_{l \in QLk} C_{kl} Q_{kl} + \sum_{k \in QK} \sum_{n \in Fk} \sum_h \sum_{i \in QIk} c_{kni} QX_{knhi} \quad (34)$$

Subjec to

$$\sum_{k \in QK} \sum_{n \in Fk} \sum_{i \in QIk} a_{kniht} QX_{knhi} - \sum_{k \in QK} \sum_{n \in Fk} \sum_{i \in QT1knht} QU_{kniht}$$

5

$$- \sum_{k \in QK} \sum_{n \in Fk} \sum_{i \in QT2knht} (QW_{kniht} + QW_{kniht(t-1)})$$

$$- \sum_{k \in QK} \sum_{n \in Fk} \sum_{i \in QT3knht} QV_{kniht} \geq b_{ht}$$

10

$$, t \in T_h, h = 1, \dots, 7, \quad (35)$$

$$QX_{knhi} = \sum_{t \in QB1knih} QU_{kniht}, \quad n \in F_k, i \in QI_k, k \in QK, h = 1, \dots, 7, \quad (36)$$

$$QX_{knhi} = \sum_{t \in QB2knih} QW_{kniht}, \quad n \in F_k, i \in QI_k, k \in QK, h = 1, \dots, 7, \quad (37)$$

15

$$QX_{knhi} = \sum_{t \in QB3knih} QV_{kniht}, \quad n \in F_k, i \in QI_k, k \in QK, h = 1, \dots, 7, \quad (38)$$

$$\sum_{l \in QLk} A_{klh} Q_{kl} = \sum_{n \in Fk} \sum_{i \in Ik} QX_{knhi}, \quad k \in QK, h = 1, \dots, 7, \quad (39)$$

20

$$\sum_{l \in QLk} Q_{kl} \leq QD_k^{\max}, \quad k \in QK, \quad (40)$$

$$QX_{knhi}, Q_{kl}, QU_{kniht}, QW_{kniht}, \text{ and } QV_{kniht} \geq 0 \text{ and integer for all } k, n, i, h, \text{ and } t. \quad (41)$$

25

Expression (34) above is the objective function that minimizes the total “fixed” cost of assigning agents to tours with various feasible work patterns plus the total wages

paid to agents. Constraint (35) ensures that the total number of agents assigned to various tours and shift lengths, and available in period t on day h minus the number of agents who are at work on day h but taking their first relief (second term on the left side of (35)), lunch (third term on the left side of (35)), and second relief (four term on the left side of (35)) in period t on day h is sufficient to meet the agent requirements in that period. Break constraints (36-38) ensure that all required breaks are scheduled for each agent who is assigned to tour k . These constraints as well as the break variables are defined only for valid breaks for a shift length. Constraint (39) ensures that the number of shifts scheduled on day h is equal to the number of agents who work on that day based on the scheduled work patterns. Constraint (40) restricts the total number of agents assigned to different feasible work patterns of tour type k to be less than or equal to the total number of agents available to assign to this tour. This constraint can easily be modified to limit the number of agents that can be assigned to a set of tours. Finally, constraint (41) restricts the work pattern, daily shift, and break variables to nonnegative integers to avoid negative values and/or fractional agent schedules.

Extensions of MILP Model (34-41)

Objective function (34) can be easily reformulated to minimize the total amount of agent time scheduled (or paid agent time) by setting C_{kl} equal to scheduled time (or paid agent time) per agent assigned to tour k and work pattern l , $l \in QL_k$, and $k \in QK$. Similarly, if agents rate their most preferred, second most preferred, etc., tours using a rating system, these ratings for various tours and work patterns can be aggregated and used in the objective value to maximize agent preference in scheduling tours and work patterns. In order to not to schedule redundant tours and increase agent costs unnecessarily, rating system should assign the lowest value to the most preferred tour, and next value higher to the second most preferred tour, etc. Consequently, the objective function will remain as of minimization type to the maximize agent preference.

The MILP model can be easily modified to develop optimal schedules when there are not enough agents to meet the agent requirements in all periods (i.e. when there are going to be agent shortages in some periods). In this case, objective function (34) is modified as follows:

$$\begin{aligned} \text{Minimize } & \sum_{i \in Q_{Lk}} \sum_{k \in Q_K} C_{kl} Q_{kl} + \sum_{k \in Q_K} \sum_{n \in F_k} \sum_h \sum_{i \in Q_{lk}} C_{kni} QX_{kni} \\ & + \sum_{t \in QT_h} \sum_h P_{ht} S_{ht} \end{aligned} \quad (42)$$

Constraint (35) is then formulated by adding O_{ht} for overstaffing and S_{ht} for shortage to its left hand side as follows:

$$\begin{aligned} & \sum_{k \in Q_K} \sum_{n \in F_k} \sum_{i \in Q_{lk}} a_{kniht} QX_{kni} - \sum_{k \in Q_K} \sum_{n \in F_k} \sum_{i \in QT1_{kniht}} QU_{kniht} \\ & - \sum_{k \in Q_K} \sum_{n \in F_k} \sum_{i \in QT2_{kniht}} (QW_{kniht} + QW_{kniht(t-1)}) \\ & - \sum_{k \in Q_K} \sum_{n \in F_k} \sum_{i \in QT3_{kniht}} QV_{kniht} + S_{ht} - O_{ht} \geq b_{ht} \end{aligned} \quad (43)$$

, $t \in T_h$, $h = 1, \dots, 7$,

$$S_{ht}, O_{ht} \geq 0, \text{ for all } t \in QT_h \text{ and } h. \quad (44)$$

If a certain amount of overstaffing is needed for other tasks (e.g. to answer email or fax based contacts), it can be assured by restricting the overstaffing to be greater than or equal to the desired level f_{ht} .

$$\sum_{t \in QTh} O_{ht} \geq f_{ht}, \quad t \in T_h, h=1, \dots, 7, \quad (45)$$

If up to a certain amount of understaffing staffing is tolerable (e.g. to see if the
 5 total cost can be reduced), this can be achieved by adding a similar constraint:

$$\sum_{t \in QTh} S_{ht} \geq d_{ht}, \quad t \in T_h, h=1, \dots, 7, \quad (46)$$

where d_{ht} is the minimum overtime desired on day h . Note that (45) and (46) can easily
 10 modified to impose limits for a specific period(s) on day h .

This case is similar to one of the cases discussed earlier for formulation (1-8).
 Constraints (2-5) and break variables in (1-8) have the same structure with constraints
 (35-38) and break variables. In both cases, the number of daily breaks scheduled is equal
 15 to the number of agents assigned to a shift and working that day. Thus, the existence of
 non-consistent daily start times and shift lengths doesn't affect the way breaks are
 scheduled in the model. Consequently, tours with more breaks or longer breaks can be
 modeled as disclosed before for formulation (1-8).

20 Since the valid work patterns are determined in advance in formulating (34-41),
 the minimum, maximum, and exact number of consecutive days off and/or total number
 of days off requirements for a tour type can be easily included in formulation (34-41).
 All possible work patterns are shown in FIG. 3. All feasible work patterns satisfying the
 days-off rules for tour k , $k \in QK$, are then identified from this list and included in QL_k .
 25 Their coefficients A_{klh} are included in constraint (39).

If a tour type has fixed days off for all agents assigned to it then all feasible work
 patterns selected for that tour should have a non-work day on that day. To reduce the size
 of the formulation, shift and break variables for the fixed days off as well as associated
 30 break constraints (36-38) are excluded from formulation (34-41).

If working hours are shorter on certain days for a tour or a contact center, the present invention extends the MILP model by excluding the daily shift variables that has a late start time resulting an end time after the early closure time for that tour or center. Break variables as well as break constraints (34-41) associated with these shift variables are excluded.

**Workforce Scheduling with Tours Requiring Both Consistent
Daily Start Times and Shift Lengths, and Non-Consistent
Start Times and Shift lengths in
a Single Skill Environment**

A workforce scheduling environment may include tours with and without consistent daily start time and shift length requirements discussed earlier. The present invention extends the MILP model to this type of workforce scheduling environment in two ways. First, even though formulation (34-41) is disclosed earlier for scheduling environments involving only tours not requiring consistent daily start times and shift length, it is extended to model such tours with a slight modification as follows; If a tour requires consistent daily start times and shift lengths, then each daily start time for this tour type represents a pseudo-tour type that has the same start time and shift length on all days. Thus, a tour that can start every 30 minutes in a 7-day-a-week-and-24-hour-a-day operation will have 48 (for 48 daily start time) pseudo tour types. Each pseudo tour k has only one daily start time in the daily start time set QI_k . Consequently, formulation (34-41) will have work pattern variables Q_{kl} , and one daily shift variable QX_{khi} , and the associated daily break variables for that shifts for tour k , and day $h = 1, \dots, 7$, for each such pseudo tour.

Moreover, if a tour has multiple shift lengths but doesn't allow varying daily shift start times, the present invention extends the MILP model by introducing pseudo tour types (one per daily start type) and formulating the shift and work pattern variables as in

formulation (34-41) with daily shift lengths in F_k . This way both consistent daily start time requirement and availability of multiple daily shift lengths are included in the MILP model.

5 Another way that the present invention extends the MILP model is to merge formulations (1-8) and (34-41) for this type of scheduling environments to obtain optimal schedules. To distinguish cost parameters, Cx_{ki} is introduced as the cost of assigning an agent to X_{ki} . Using the definitions of the parameters and variables given for formulations (1-8) and (34-41), merging the objective functions (1) and (34) and demand
10 constraints (2) and (35), and including the necessary break and days-off constraints from formulations (1-8) and (34-41), the present invention extends the MILP model for a scheduling environment involving both consistent and non-consistent shift start times and shift lengths as follows:

$$15 \text{ Minimize } \sum_{k \in K} \sum_{i \in I_k} Cx_{ki} X_{ki} + \sum_{k \in QK} \sum_{l \in Q_k} C_{kl} Q_{kl} + \sum_{k \in QK} \sum_{n \in F_k} \sum_h \sum_{i \in Q_k} c_{kni} QX_{kni} \quad (47)$$

Subject to

$$20 \begin{aligned} & \sum_{k \in K} \sum_{i \in I_k} a_{kiht} X_{ki} - \sum_{k \in K} \sum_{i \in I_k} a_{kiht} (Y_{kih} + Y_{ki(h-1)}) - \sum_{k \in K} \sum_{i \in T1kht} U_{kiht} \\ & - \sum_{k \in K} \sum_{i \in T2kht} (W_{kiht} + W_{kih(t-1)}) - \sum_{k \in K} \sum_{i \in T3kht} V_{kiht} \\ & + \sum_{k \in QK} \sum_{n \in F_k} \sum_{i \in Q_k} a_{kniht} QX_{kni} - \sum_{k \in QK} \sum_{n \in F_k} \sum_{i \in QT1kht} QU_{kniht} \\ & - \sum_{k \in QK} \sum_{n \in F_k} \sum_{i \in QT2kht} (QW_{kniht} + QW_{kniht(t-1)}) \\ 25 & - \sum_{k \in QK} \sum_{n \in F_k} \sum_{i \in QT3kht} QV_{kniht} \geq b_{ht} \end{aligned}$$

$$t \in T_h, h = 1, \dots, 7 \quad (48)$$

$$X_{ki} - Y_{kih} - Y_{ki(h-1)} = \sum_{t \in B1kih} U_{kiht} \quad , i \in I_k, k \in K, h = 1, \dots, 7, \quad (49)$$

$$X_{ki} - Y_{kih} - Y_{ki(h-1)} = \sum_{t \in B2kih} W_{kiht} \quad , i \in I_k, k \in K, h = 1, \dots, 7, \quad (50)$$

$$5 \quad X_{ki} - Y_{kih} - Y_{ki(h-1)} = \sum_{t \in B3kih} V_{kiht} \quad , i \in I_k, k \in K, h = 1, \dots, 7, \quad (51)$$

$$X_{ki} = \sum_h Y_{kih} \quad , i \in I_k, k \in K, \quad (52)$$

$$10 \quad QX_{knhi} = \sum_{t \in QB1kniht} QU_{kniht} \quad , n \in F_k, i \in QI_k, k \in QK, h = 1, \dots, 7, \quad (53)$$

$$QX_{knhi} = \sum_{t \in QB2kniht} QW_{kniht} \quad , n \in F_k, i \in QI_k, k \in QK, h = 1, \dots, 7, \quad (54)$$

$$QX_{knhi} = \sum_{t \in QB3kniht} QV_{kniht} \quad , n \in F_k, i \in QI_k, k \in QK, h = 1, \dots, 7, \quad (55)$$

$$15 \quad \sum_{l \in QLk} A_{klh} Q_{kl} = \sum_{n \in F_k} \sum_{i \in I_k} QX_{knhi} \quad , k \in QK, h = 1, \dots, 7, \quad (56)$$

$$\sum_{i \in I_k} X_{ki} \leq D_k^{\max} \quad , k \in K, \quad (57)$$

$$\sum_{l \in QLk} Q_{kl} \leq QD_k^{\max} \quad , k \in QK, \quad (58)$$

$$20 \quad X_{ki}, Y_{kih}, QX_{knhi}, Q_{kl}, U_{kiht}, W_{kiht}, \text{ and } V_{kiht}, QU_{kniht}, QW_{kniht}, \text{ and } QV_{kniht} \geq 0 \text{ and integer for all } k, n, i, h, \text{ and } t, \quad (59)$$

where objective function (47) combines objectives functions (1) and (34), constraint (48) combines agent, break, and days-off variables specifying agent availability in a period from constraints (2) and (35), constraints (49-51) are the break balance constraints for the tours requiring consistent daily shift start times and shift lengths, constraint (52) ensures that sufficient number of days-off are scheduled for the agents assigned to tours requiring consistent daily shift start times and shift lengths, constraints (53-55) ensure that sufficient number of first, lunch, and second relief breaks are scheduled for the agents assigned to tours not requiring consistent daily shift start times and shift lengths, constraint (56) ensures that the number of shifts scheduled on a given day is equal to the number agents who are assigned to work on that day by the work patterns scheduled, constraints (57) and (58) ensure that the numbers of agents assigned to various tours requiring consistent or non-consistent daily shift start times and shift lengths don't exceed the limits on the agent availability for these tours. Finally, constraint (59) restricts the tour, days-off, daily shift, work pattern, and break variables to nonnegative integers to avoid negative values and/or fractional agent schedules.

Depending on the scheduling environment and its tour and shift requirements, the present invention extends the MILP model by merging formulations (1-8), (14-23), (24-31), (34-41) or any extensions of these as disclosed before in a manner similar to formulation (47-59) above. It should be clear to those skilled in the art that the specific embodiments and the MILP model disclosed above can be readily utilized as a basis and extended to other scheduling environments involving the types of tours considered in formulations (1-8), (14-23), (24-31), and (34-41), and their variations without departing from the scope of the invention.

Workforce Scheduling in a Skills Based Environment

The present invention extends the MILP model to scheduling environments involving multiple agent skill groups, agent work groups, and multiple contact types requiring different agent skills. An agent skill group contains agents with the same skill

sets. Let R be the set of different contact types with known agent requirements in each period t and day h . Let the number of agents with qualified skills for contact type $r \in R$ required in period t on day h be b_{ht}^r . Suppose now that there are a number of agent skill groups $j \in J$, each with a unique set of skills. Agent requirements should reflect the multi-skill efficiency expected in a skills-based routing environment due to agents with multiple skills and, depending on the scheduling environment, may be specified for skills groups rather than contact types. Let the set of contact types that skills group j is qualified to provide service be N_j , $j \in J$. A contact type may be served by multiple agent groups. Let the set of skills groups that can service contact type r be M^r .

The present invention extends the MILP model for skills based scheduling by formulating the agent assignment to contact types, and contact type based agent requirements fulfillment in separate constraints. To facilitate the description, consider formulation (34-41) and constraint (35). Further, consider the availability of agents from group j . The number of agents that are available from group j in period t on day h is equal to the number of agents scheduled to work on day h minus the number of agents taking a relief or lunch break in period t . Let G_{ht}^{jr} be the number of agents from skill group j allocated to serve type r contacts, where $r \in N_j$, and $j \in M^r$. These variables are also referred to as “skill allocation variables”. Thus,

$$\sum_{k \in QKj} \sum_{n \in Fk} \sum_{i \in Qlk} a_{kniht} QX_{kniht}^j - \sum_{k \in QKj} \sum_{n \in Fk} \sum_{i \in QT1kniht} QU_{kniht}^j \\ - \sum_{k \in QKj} \sum_{n \in Fk} \sum_{i \in QT2kniht} (QW_{kniht}^j + QW_{kniht(t-1)}^j)$$

$$- \sum_{k \in QKj} \sum_{n \in Fk} \sum_{i \in QT3kniht} QV_{kniht}^j - \sum_{r \in N_j} G_{ht}^{jr} = 0,$$

$$, j \in J, t \in T_h, h = 1, \dots, 7, \quad (61)$$

The first term on the left side of (61) is the number of agents from group j scheduled in period t on day h . The second through fourth terms on the left side are the numbers of

agents who are scheduled to be on a relief or lunch break during this period. Thus, the difference is equal to the number of agents who are scheduled and working (available) during period t on day h . Finally, the fifth term is the sum the numbers of agents allocated to various contact types in N_j .

5

The present invention extends the MILP model into the skills based scheduling environment in two different ways. First approach involves merging formulations (14-23) and (34-41). Consider constraints (15) and (35). In the skills based scheduling environment, the right sides of these two constraints are merged as follows:

10

$$\begin{aligned}
 f_{ht}^j(X, Y, Z, U, W, V, QX, Q, QU, QW, QV) = & \sum_{k \in K_j} \sum_{i \in I_k} a_{kiht} X_{ki}^j \\
 & - \sum_{k \in K_j} \sum_{i \in I_k} a_{kiht} (Y_{kih}^j + Y_{ki(h-1)}^j) - \sum_{k \in K_j} \sum_{i \in I_k} \sum_{m \neq h, (h-1)} a_{kiht} Z_{kimh}^j \\
 & - \sum_{k \in K_j} \sum_{i \in T1kht} U_{kiht}^j - \sum_{k \in K_j} \sum_{i \in T2kht} (W_{kiht}^j + W_{kih(t-1)}^j) \\
 & - \sum_{k \in K_j} \sum_{i \in T3kht} V_{kiht}^j \\
 & + \sum_{k \in QK_j} \sum_{n \in F_k} \sum_{i \in Qlk} a_{kniht} QX_{kni}^j \\
 & - \sum_{k \in QK_j} \sum_{n \in F_k} \sum_{i \in QT1kniht} QU_{kniht}^j \\
 & - \sum_{k \in QK_j} \sum_{n \in F_k} \sum_{i \in QT2kniht} (QW_{kniht}^j + QW_{kniht(t-1)}^j) \\
 & - \sum_{k \in QK_j} \sum_{n \in F_k} \sum_{i \in QT3kniht} QV_{kniht}^j
 \end{aligned}$$

15

20

$$, j \in J, t \in T_h, h = 1, \dots, 7, \quad (62)$$

The terms on the right hand side of (62) represented by $f_{ht}^j(X, Y, Z, U, W, V, QX, Q, QU, QW, QV)$ give the number of agents from group j assigned to various tours (with and without consistent start time and length) and are available in period t on day h . The notation introduced for (1-61) is now extended to distinguish agent skills groups. For example, $X = \{ X_{ki}^j : i \in I_k, k \in K_j, h = 1, \dots, 7 \}$ where X_{ki}^j is as defined before except it now

25

represents the number of agents from group j assigned to tour k with a daily start time of i . The sets $Y, Z, U, W, V, QX, Q, QU, QW$, and QV are defined similar to the set X to include the respective variables for the agent groups in J . Constraint (61) is now extended to the skills based scheduling MILP model of the present invention as follows.

5

$$f_{ht}^j(X, Y, Z, U, W, V, QX, Q, QU, QW, QV) - \sum_{r \in N_j} G_{ht}^{jr} = 0, \\ , j = 1, \dots, n, t \in T_h, h = 1, \dots, 7, \quad (63)$$

10 Given the agent allocation G_{ht}^{jr} in every period and day for different contact types, agent requirement constraints for contact type r are formulated in the MILP model as follows.

$$\sum_{j \in M_r} e^{jr} G_{ht}^{jr} \geq b_{ht}^r, \quad , r \in R, t \in T_h, h = 1, \dots, 7, \quad (64)$$

15

where e^{jr} is the relative efficiency of an agent from agent group j in serving contact type r with respect to an agent whose primary skill (highest proficiency level) is serving contact type r , $e^{jr} \in [0, 1]$. For instance, if $e^{jr} = 1$, agent group j is equally efficient, and if $e^{jr} = 0.75$, agents in group j are 75% efficient with respect to an agent whose primary skill is serving contact type r . Values of G_{ht}^{jr} variables do not have to be integer.

20

Variable definitions presented before are modified to include agent group $j \in J$ as a superscript in all variables. To schedule breaks and days off for different agent groups, constraints (16-18) and (36-38), except for variable definitions for different agent skill groups, remain the same since the agent allocation does not affect the break and days-off scheduling constraints. Objective functions (14) and (34) are merged and modified to include potential wage differentials for different agent groups. The MILP model of the present invention is now disclosed below.

25

$$\begin{aligned} \text{Minimize } \sum_{j \in J} \sum_{i \in I_k} \sum_{k \in K_j} C_{ki}^j X_{ki}^j + \sum_{j \in J} \sum_{l \in Q_{Lk}} \sum_{k \in Q_{Kj}} C_{kl}^j Q_{kl}^j \\ + \sum_{j \in J} \sum_{k \in Q_{Kj}} \sum_{n \in F_k} \sum_h \sum_{i \in Q_{Ik}} C_{kni}^j QX_{knh}^j \end{aligned} \quad (65)$$

5 Subject to

$$\sum_{j \in M_r} e^{jr} G_{ht}^{jr} \geq b_{ht}^{r}, \quad r \in R, \quad t \in T_h, \quad h = 1, \dots, 7, \quad (66)$$

$$\begin{aligned} f_{ht}^j(X, Y, Z, U, W, V, QX, Q, QU, QW, QV) \\ - \sum_{r \in N_j} G_{ht}^{jr} = 0 \end{aligned}$$

10

$$, j \in J, \quad t \in T_h, \quad h = 1, \dots, 7, \quad (67)$$

$$X_{ki}^j - Y_{kih}^j - Y_{ki(h-1)}^j - \sum_{m \neq h, (h-1)} Z_{kimh}^j = \sum_{t \in B1kih} U_{kiht}^j, \quad j \in J, \quad i \in I_k, \quad k \in K_j, \quad h = 1, \dots, 7, \quad (68)$$

$$15 \quad X_{ki}^j - Y_{kih}^j - Y_{ki(h-1)}^j - \sum_{m \neq h, (h-1)} Z_{kimh}^j = \sum_{t \in B2kih} W_{kiht}^j, \quad j \in J, \quad i \in I_k, \quad k \in K_j, \quad h = 1, \dots, 7, \quad (69)$$

$$X_{ki}^j - Y_{kih}^j - Y_{ki(h-1)}^j - \sum_{m \neq h, (h-1)} Z_{kimh}^j = \sum_{t \in B3kih} V_{kiht}^j, \quad j \in J, \quad i \in I_k, \quad k \in K_j, \quad h = 1, \dots, 7, \quad (70)$$

$$r_k Y_{kih}^j = \sum_{l \neq h, (h+1)} Z_{kihl}^j, \quad j \in J, \quad i \in I_k, \quad k \in K_j, \quad h = 1, \dots, 7, \quad (71)$$

20

$$\begin{aligned} Z_{kihm}^j \leq Y_{kih}^j, \quad j \in J, \quad i \in I_k, \quad k \in K_j, \quad h = 1, \dots, 7, \\ m \neq h, (h+1) \end{aligned} \quad (72)$$

$$X_{ki}^j = \sum_h Y_{kih}^j, \quad j \in J, \quad i \in I_k, \quad k \in K_j, \quad (73)$$

25

$$QX_{knh}^j = \sum_{t \in Q_{B1knh}} QU_{knh}^j, \quad j \in J, \quad n \in F_k, \quad i \in Q_{Ik}, \quad k \in Q_{Kj}, \quad h = 1, \dots, 7, \quad (74)$$

$$QX_{knh}^j = \sum_{t \in QB2knh} QW_{kniht}^j, \quad j \in J, n \in F_k, i \in QI_k, k \in QK_j, h = 1, \dots, 7, \quad (75)$$

$$QX_{knh}^j = \sum_{t \in QB3knh} QV_{kniht}^j, \quad j \in J, n \in F_k, i \in QI_k, k \in QK_j, h = 1, \dots, 7, \quad (76)$$

$$5 \quad \sum_{l \in QL_k} A_{klh} Q_{kl}^j = \sum_{n \in F_k} \sum_{i \in I_k} QX_{knh}^j, \quad j \in J, k \in QK_j, h = 1, \dots, 7, \quad (77)$$

$$\sum_{i \in I_k} X_{ki}^j \leq D_k^{j \max}, \quad j \in J, k \in K_j, \quad (78)$$

$$\sum_{l \in QL_k} Q_{kl}^j \leq QD_k^{j \max}, \quad j \in J, k \in QK_j, \quad (79)$$

10

$$X_{ki}^j, Y_{kih}^j, Z_{kihm}^j, U_{kiht}^j, W_{kiht}^j, V_{kiht}^j, QX_{knh}^j, Q_{kl}^j, QU_{kniht}^j, QW_{kniht}^j, QV_{kniht}^j \geq 0 \text{ and} \\ \text{integer for all } j, k, n, i, h, \text{ and } t, \text{ and } G_{ht}^r \geq 0 \text{ for all } j, r, h, \text{ and } t. \quad (80)$$

15 where $f_{ht}^j(X, Y, Z, U, W, V, QX, Q, QU, QW, QV)$ is as defined in equation (62) above.

G_{ht}^r in formulation (65-80) above doesn't have to be integer even though it represents the number of agents from group j allocated to serve type r contacts. Hence, formulation (65-80) is a Mixed Integer Programming (MILP) formulation involving both
20 integer and continuous variables.

Note that MILP formulation (65-80) may also be modified to have multiple daily shift lengths by modifying f_{ht}^j to include the set of shift lengths F_k , and appropriate a_{kiht} coefficients for $X_{ki}^j, i \in I, k \in K_j$, in (62).

25

In order to prevent the notation becoming more taxing, the MILP model of the present invention for optimal workforce scheduling in a skills based scheduling environment is presented for a combination of the scheduling environments disclosed before for formulations (15-23) and (34-41). Depending on the scheduling environment

and its tour and shift requirements, the present invention extends the MILP model by merging formulations (1-8), (14-23), (24-31), (34-41) or any extension of these as disclosed before in a manner similar to the MILP model presented above. It should be clear to those skilled in the art that the specific embodiments and the MILP model disclosed above can be readily utilized as a basis and extended to other scheduling environments without departing from the scope of the invention.

The present invention also extends the MILP model to obtain a second formulation by defining pseudo tours for the tours requiring consistent daily shift start times as follows. Consider a tour type, say tour k , $k \in K_j$, available for skills group j . Set K_j contains the valid tour types for skills group j that require consistent daily shift start times. Constraints (68-73), (78), and six terms on the right side of (62) involving variables X_{ki}^j , Y_{kih}^j , Z_{kih}^j , U_{kiht}^j , W_{kiht}^j , and V_{kiht}^j specify the break placement and days off scheduling requirements and agent availability for tours in K_j . The present invention extends the MILP model for the second formulation by first considering all valid work patterns given in FIG. 3 and identifying the patterns satisfying constraints (71-73) for $k \in K_j$. A distinct pseudo tour is then defined for every daily start time specified in I_k , $k \in K_j$, with the valid work patterns identified. Let one such pseudo tour be α . Then, α has only one daily start time in OI_α representing one of the daily start times in I_k , has the same break placement and days off scheduling requirements with tour k , $k \in K_j$, and has the work patterns in OL_α . After these modifications, pseudo tours defined this way are treated like the tours not requiring consistent daily start times and included in the tours set QK_j available for agent skill group j . Since OI_α has only one daily start time, this satisfies the consistent daily shift start time requirement. Also, for tour α , F_α may have more than one shift length specified for tour k , $k \in K_j$. Another way the present invention extends the MILP model is by merging notation used in constraints (68-70) and (74-76), and constraints (78) and (79), and removing the redundant notation and constraints as follows.

$$\text{Minimize } \sum_{j \in J} \sum_{l \in QLk} \sum_{k \in QKj} C_{kl}^j Q_{kl}^j + \sum_{j \in J} \sum_{k \in QKj} \sum_{n \in Fk} \sum_h \sum_{i \in Qlk} c_{kni}^j QX_{knh}^j \quad (81)$$

Subject to

$$\sum_{j \in Mr} e^{jr} G_{ht}^{jr} \geq b_{ht}^r, \quad r \in R, t \in T_h, h = 1, \dots, 7, \quad (82)$$

5

$$f_{ht}^j(QX, Q, QU, QW, QV) - \sum_{r \in Nj} G_{ht}^{jr} = 0, \quad j \in J, t \in T_h, h = 1, \dots, 7, \quad (83)$$

$$QX_{knh}^j = \sum_{t \in QB1knh} QU_{kniht}^j, \quad j \in J, n \in Fk, i \in QI_k, k \in QK_j, h = 1, \dots, 7, \quad (84)$$

$$10 \quad QX_{knh}^j = \sum_{t \in QB2knh} QW_{kniht}^j, \quad j \in J, n \in Fk, i \in QI_k, k \in QK_j, h = 1, \dots, 7, \quad (85)$$

$$QX_{knh}^j = \sum_{t \in QB3knh} QV_{kniht}^j, \quad j \in J, n \in Fk, i \in QI_k, k \in QK_j, h = 1, \dots, 7, \quad (86)$$

$$\sum_{l \in QLk} A_{klh} Q_{kl}^j = \sum_{n \in Fk} \sum_{i \in Qlk} QX_{knh}^j, \quad j \in J, k \in QK_j, h = 1, \dots, 7, \quad (87)$$

15

$$\sum_{l \in QLk} Q_{kl}^j \leq QD_k^{j \max}, \quad j \in J, k \in QK_j, \quad (88)$$

$QX_{knh}^j, Q_{kl}^j, QU_{kniht}^j, QW_{kniht}^j$, and $QV_{kniht}^j \geq 0$ and integer for all j, k, n, i, h, t , and $G_{ht}^{jr} \geq 0$ for all j, r, h , and t . (89)

20

where $f_{ht}^j(X, U, W, V, QX, Y, Z, Q, QU, QW, QV)$ is now expressed as given below.

$$\begin{aligned} f_{ht}^j(QX, Q, QU, QW, QV) = & \sum_{k \in QKj} \sum_{n \in Fk} \sum_{i \in Qlk} a_{kniht} QX_{knh}^j \\ & - \sum_{k \in QKj} \sum_{n \in Fk} \sum_{i \in QT1knh} QU_{kniht}^j \\ & - \sum_{k \in QKj} \sum_{n \in Fk} \sum_{i \in QT2knh} (QW_{kniht}^j + QW_{kniht(t-1)}^j) \end{aligned}$$

25

$$- \sum_{k \in QKj} \sum_{n \in Fk} \sum_{i \in QT3knht} QV_{kniht}^j$$

$$, j \in J, t \in T_h, h = 1, \dots, 7, \quad (90)$$

5 Extension of formulations (65-80) and (81-89) to the cases discussed in relation to
formulations (1-8) and (34-41) is straightforward. It should be clear to those skilled in
the art that the information disclosed in relation to formulations (1-8) and (34-41) can be
readily utilized as a basis to extend formulations (65-80) and (81-89) of the present
invention to the cases involving (i) minimizing scheduled/paid time or maximizing agent
10 preference, (ii) schedule generation when there aren't enough agents (i.e. there are agent
shortages in some periods for some contact types), (iii) tours with longer and/or more/less
breaks, (iv) fixed days-off, and (v) days with a closure time earlier than the other days.
Accordingly, it is to be understood that the embodiments of the invention herein
described are merely illustrative of the application of the principles of the invention.

15 Reference herein to details of the illustrated embodiments are not intended to limit the
scope of the claims, which themselves recite those features regarded as essential to the
invention.

20 **Description of the Method for finding an Optimal Solution of the MILP model and an Optimal Schedule in a Skills Based Scheduling Environment**

Formulations (65-80), and (81-89), and their extensions disclosed before can be
25 solved to obtain a global optimal solution in a number of ways. Such a global optimal
solution satisfies the necessary and sufficient conditions for optimality of the MILP
model of the invention. The solution method of the invention will be discussed in the
context of formulations (65-80) and (81-90). However, the solution method of the
present invention can also be used to solve optimally formulation (1-8) and its extensions,
30 and formulation (24-31) and its extensions as well as formulation (47-59) disclosed

before. Therefore, the scope of the solution method of the invention also includes such MILP models.

The branch and cut (B&C) algorithm is a well known technique for optimally solving MILP formulations such as (65-80), and (81-89) (Wolsey, 1998). A computer program implementing the B&C algorithm can be developed using any programming language or can be obtained from a number of software companies including Ilog, Inc., and Dash Optimization. These providers are also offering their B&C algorithm in the form of a callable library that can be called from an integrated MILP model generator and optimizer module. In either case, the B&C algorithm forms a part of the solution method of the present invention.

The MILP formulation generator ((3) in FIG. 1) of the invention is implemented in a computer program. Given the agent and skills requirements for each planning period $t \in QT_h$, $h = 1, \dots, 7$, and center information regarding agent groups, tours and shift parameters, the MILP formulation generation program develops a formulation (step (11)) of the scheduling problem using the MILP formulations and their extensions (1-89) of the invention disclosed before. To generate a formulation, the MILP formulation generation program calculates the coefficients of various shift, break, days-off, work pattern, overage and underage, agent allocation variables in the respective equations and inequalities of the MILP model of the invention for the scheduling environment considered. This information is used by the MILP formulation generator program to generate data arrays containing the coefficients of the decision variables in every constraint defined for the particular scheduling environment to meet its requirements. These arrays are then passed to the optimizer module (step (12)). It is to be understood that a number of versions of the MILP formulation generation method can be developed for the formulations, and their extensions disclosed above without departing from the scope of the invention.

The optimizer module is a computer program working with the MILP formulation generator program, and uses a B&C algorithm supplemented by the RA algorithm of the

invention. The purpose of the RA algorithm is to search for integer feasible solutions for the MILP model generated together with the B&C algorithm. If a sub-problem is infeasible (i.e. the sub-problem has no integer or non-integer solution), or has an integer feasible solution (that is, a solution satisfying all constraints of the model including the integrality constraints for all decision variables), or has a solution that is inferior to the best available lower bound for the optimal integer solution of the MILP model, the B&C algorithm doesn't call the RA algorithm since it can't develop a new or different integer solution. Otherwise, the B&C algorithm calls the RA algorithm after obtaining a non-integer but feasible solution to a node. At each such node (or after every any number of such nodes), using the non-integer solution found by the B&C algorithm to that node, the RA algorithm makes an attempt to find an integer feasible solution to the MILP model of the invention while trying to achieve the best objective value.

The B&C algorithm first solves the continuous version (i.e. Linear Programming (LP) relaxation) of an MILP model (Wolsey, 1998) (step (17)). This node is called node zero (subsequent nodes are numbered sequentially). If the continuous solution thus found is integer feasible and optimal (step (20)), then the optimizer terminates with an optimal solution to the MILP model (step (21)). If this solution is not integer feasible (i.e. if some integer restricted variables have non-integer values), then the B&C algorithm starts creating nodes (sub-problems) by adding constraints on certain decision variables to eliminate non-integer values. In this case, the objective value of the solution of the LP relaxation is a lower bound on the lowest objective value that can ever be achieved for the scheduling environment formulated in the MILP model. That is, no solution to the MILP model can have an objective value better than the lower bound thus obtained. After solving the LP relaxation at node 0, if the solution is a feasible solution to the LP relaxation but violates some integrality constraints (step (22)), the B&C algorithm calls the RA algorithm. If the RA algorithm locates an integer feasible solution, its objective value is saved as the best heuristic objective value, and the integer solution is returned to the B&C algorithm (step (100A)) as the best-known integer solution (if no integer solution is available beforehand). In the subsequent nodes, the RA algorithm of the invention proceeds as outlined below.

Step 23: Obtain the objective value for the best integer solution known from the B&C algorithm. If this value is better than the best heuristic objective value in the RA algorithm, update the best heuristic objective value.

5

Step 24: Obtain the values of the shift, breaks, days-off, work pattern, and skill allocation variables (X, U, W, V, Y, QX, Q, QU, QW, QV) in the non-integer solution to the node solved from the B&C algorithm.

10 Step 25: Round the fractional values of all U, W, V, Y, QX, QU, QW, QV, and G variables down (i.e. truncate the fractional values). As for X and Q variables, round these variables down by their fractional parts are less than or equal to 0.50. Keep a list of the variables with a fractional part greater than 0.50. Find the total number of agents assigned to each tour type. Round the variables that had a
15 fractional part greater than 0.50 up, one at a time, if the new value is not violating the maximum agent constraints (e.g. constraints (78-79)). If the maximum agent limit is reached for a tour group, round the remaining variables down even if their fractional parts are greater than 0.50. Compute the agent shortages and excesses in each period for all contact types using the rounded values of decision variables.

20

Step 26: Compare the days-off variables Y and Z with shift variables X for tours with consistent daily start time and shift length, and daily shift variables QX with work pattern variables Q. If additional days off or daily shifts are needed to satisfy tour requirements, go to step (27). Otherwise, go to (28).

25

Step 27: Schedule more days off (i.e. increase the values of appropriate days-off variables) for the tours with consistent start times on the days with the lowest maximum per period, or total shortages during a tour's shift span to satisfy days off requirements for every agent currently scheduled. Similarly, schedule daily
30 shifts to have sufficient number of daily shifts scheduled for the tours not requiring consistent daily shift start times on the days and shift start times with the

highest maximum per period shortage, or total shortage covered by that shift.

Thus, the tour requirements for days off, and daily shifts (e.g. constraints (71-73) and (77)) will be satisfied by these variables.

5 Step 28: Compare the break variables, U, W, and V with the shift variables X and days-off variables Y, and QX, QU, and QW with daily shift variables QX. If the required number of breaks is scheduled for every tour, go to step (30). Otherwise, go to step (29).

10 Step 29: If more breaks are scheduled on a day for a tour, unschedule breaks (i.e. decrease the value of appropriate break variables) to have the correct number of breaks scheduled. If fewer than the required number of breaks are scheduled, schedule additional breaks in the periods (contagious if a break is more than one period long within the break window of the respective break) that has the lowest
15 maximum per period shortage, or total shortage (or if no period has shortage, then the highest maximum per period excess, or total excess) to satisfy the break balance requirements (e.g. constraints (68-70) and (74-76).

20 Step 30: Compute the agent shortages and excesses using the agent allocation variables G for each contact type, planning period, and day.

Step 31: Compute the scheduled agent availability in each period for all agent groups (i.e. $f_{ht}^j(X, U, W, V, QX, Y, Q, QU, QW, QV)$ in equation (67)). Compare the
25 number of agents available in each agent group j in a planning period, say $t \in T_h$ on day h, with the values of the agent allocation variables G associated with this agent group in planning period t. If the number of agents available to work for agent group j is not equal to the sum of the current values of the agent allocation variables (from step 25) for all contact types in N_j in period t on day h, adjust
30 values of associated allocation variables to meet the requirements by contact types in N_j . Prioritize contact types in each period by (i) maximum shortage, (ii)

whether this agent group is the only agent group (or the number of agent groups and tour groups) in M^f , the set of agent groups that can serve a contact type, (iii) lowest cost per period per agent to meet shortage for a contact type. This allocation process can be carried out in a number of ways. These allocation

5 criteria are implemented in a nested fashion so that if contact types have the same merits as far as criteria (i) is concerned, then the tie is broken by applying criteria (ii), and so on. This process results in a schedule that allocates all agents scheduled. Update period shortages and excesses.

- 10 Step 32: The schedule constructed using the non-integer solution found by the B&C algorithm for a node satisfies all constraints (e.g. formulation (65-80)), but may not satisfy the agent and skill requirements (e.g. constraint (66)). If all agent and skill requirements are satisfied in all periods to be scheduled, then go to step 39. Otherwise, go to step 33.

15

Step 33: If all available agents are scheduled, go to step 34. Otherwise, go to step 37.

Step 34: If a feasible schedule satisfying all agent requirements is found either by the B&C algorithm, or in the earlier calls to the RA algorithm, go to step 35.

20 Otherwise, go to step 36.

Step 35: Discard the current infeasible solution (that is, an integer solution with agent shortages). Stop the RA algorithm, and return control back to the B&C algorithm (step (17)).

25

Step 36: If an integer feasible solution to the MILP model hasn't been found by the B&C and RA Algorithms at earlier tries, compare the new infeasible solution with the infeasible solution recorded in earlier calls to the RA algorithm, and update the best infeasible solution if the new one has fewer period shortages. If this is node

30 zero, store the infeasible solution if there are no known solution. Stop the RA algorithm, and return back to the B&C algorithm (step (17)).

Step 37: Given the current agent allocations and requirements, find the agent requirement constraints that are not satisfied (i.e. have shortages) for all contact types and periods. Considering all agent groups $j \in J$, and tour types $k \in K_j$ and $k \in QK_j$, find a
 5 tour schedule that will reduce the highest number of period shortages per dollar, and schedule the tour with an agent. To specify the tour schedule for the new agent, find a daily shift start time, days-off schedule, and breaks for work days if the tour type requires consistent daily start time, or work pattern daily shift start times, shift lengths, and break times for work days if the tour type doesn't require
 10 consistent daily start time and shift length as follows:

37a: For each agent skill group j with available agents (i.e. the number of agents already scheduled in the rounded solution found in step (25) is less than the maximum number of agents available for an agent skill group),
 15 consider tour type k that is available for this agent group. If the tour requires consistent daily start time and shift length, that is $k \in K_j$, go to 37b. Otherwise, $k \in QK_j$ and go to 37c.

37b: Consider a daily start times in P_k for tour type k , and all contact types in R .
 20 Determine the period with the highest shortage level covered by this tour (referred to as "peak shortage coverage" in the remainder of this document). In order to determine the highest peak shortage coverage by a new agent, allocation of an agent's time should be decided based on the shortages by contact types to maximize this measure. Given the allocation
 25 of agent's time, determine also the total number of agent shortage periods (referred to as "total shortage coverage" in the remainder of this document) covered by this tour. All subsequent references to shortages and excesses refer to this allocation of the new agent's time. Consider the days-off schedule for this agent. Schedule the required number of days off
 30 on the days with the highest total excess during the tour schedule (or if all days have shortages, then the days with the lowest total shortage). Days-

off schedule should adhere to minimum number of consecutive days off as well as fixed days off requirements. Adjust the total shortages covered by this agent. Go to 37d.

- 5 **37c:** Consider a work pattern in QL_k^j for tour type k , and all contact types in R .
 For the selected work pattern, determine the daily start time for each work day that will result in the highest peak shortage coverage by this tour. In order to determine the maximum shortage coverage by a new agent, allocation of an agent's time should be decided based on the shortages by
 10 contact types to maximize this measure. Given the allocation of agent's time, also determine the total number of shortage periods covered by this tour. All subsequent references to shortages and excesses refer to these allocations. Go to 37d.

- 15 **37d:** Consider the work days for the tour identified in 37b or 37c. Schedule the relief and lunch breaks for work days into the periods within the specified break windows with the highest total excess (or, if all periods within a break window has shortages, then, then the periods with the lowest total shortage). Adjust the total shortage covered by this agent.

- 20 **37e:** If tour k requires consistent daily shift start times and lengths, compare the peak shortage covered by tours starting at different times in I_k^j and find the start time i^* with the highest peak shortage covered. If there are ties between two or more start times in I_k^j , select the one with the highest
 25 shortage coverage per dollar

$$(\text{total shortage coverage}) / C_{ki}^j$$

- 30 If tour k does not require consistent daily shift start times and lengths, compare the peak shortage covered by tours with different work patterns in QL_k^j and find the work pattern l^* with the highest peak shortage

covered. If there are ties between two or more work patterns in QL_k^j select the one with the highest shortage coverage per dollar

(total shortage coverage)

5

$$\frac{(\text{total shortage coverage})}{(C_{ki}^j + \sum_h (\text{number of hours worked on day } h) * c_{kni}^j)}$$

10

37f: If tour k for agent group j is the first tour evaluated, then store the current skills group, tour schedule, and agent allocation as the best ones found so far, and go to step 37a. Otherwise, if the peak shortage covered by tour k is higher than the highest peak shortage covered by any tour and skills group considered so far, update the best skills group, tour schedule, and agent allocation with the current ones. If there is a tie between the current tour schedule and skills group, and the best skills group and tour schedule found so far, select the one with the highest

15

$$\frac{(\text{total shortage coverage})}{C_{ki}^j}, \text{ for } i \text{ when } k \in K_j,$$

20

(total shortage coverage)

$$\frac{(\text{total shortage coverage})}{(C_{ki}^j + \sum_h (\text{number of hours worked on day } h) * c_{kni}^j)}$$

, for l when $k \in QK_j$,

25

Also update the peak shortage covered, and the total shortage coverage for the current tour schedule. Go to **37g** if all skills groups and all tours are evaluated. Otherwise, go to **37a**.

30

37g: Add the best tour schedule found in steps **37a-37f** to the schedule developed

in steps 16-37. Update the values of shift, days-off, break, and agent allocation X , Y , U , W , V , QX , Q , QU , QW , QV , and G variables. Update shortages and excesses using the expanded schedule with the new agent.

- 5 Step 38: If there are shortages in any period and contact type, go to step 33. Otherwise, a feasible schedule is found. Go to step 39.

- Step 39: Consider the agents scheduled in the expanded schedule. Start with the most expensive skills group and consider the tours scheduled for this group. If there is
10 a tour that can be removed without creating any shortages in any period and contact types, the current solution has redundant tours. Consider all skills groups in this fashion to determine redundant agents. Go to step 40 to remove the redundant tours. Otherwise, go to step 41.

- 15 Step 40: Remove all redundant tours together with the shift, break, and days-off schedules associated with them.
-

- Step 41: Compare the objective value for the current schedule with that of the best feasible solution to the MILP model found so far (if no feasible schedule is
20 known, then set the cost for best found schedule to a large value such as 10^{+15}) by both the B&C algorithm and the RA algorithm. Go to step 42 if the current solution has a better objective value than the best known feasible solution. Otherwise go to step 44.

- 25 Step 42: Update the best known feasible schedule with the current solution. Update the values received from the B&C for the variables.

Step 43: Stop the RA algorithm, and return the new solution to the B&C algorithm as the current best integer solution (step (16)).

- 30 Step 44: Discard the current solution. Stop the RA algorithm, and return to the B&C algorithm as the current best integer solution (step (17)).

The RA algorithm can also be supplemented with a two-way interchange, or general n-way interchange step so that, when an integer feasible solution is found, the RA algorithm can make an attempt to improve it by replacing one or more scheduled tours with other tours that are not currently scheduled. This is a well-known enhancement in the field of optimization that is widely applied to a number of other optimization problems. Also, if chosen, the solution method of the invention may also be terminated prior to finding an optimal solution if, for example, (i) the objective function for the best known integer solution differs less than a pre-specified percentage from the best lower bound found by the B&C algorithm, (ii) a pre-specified number of sub-problems (e.g. one or more sub-problems) are solved by the B&C and RA algorithms, or (iii) a pre-specified amount of time elapses after locating the first integer feasible solution to the MILP. A solution that is either optimal or satisfying these conditions is referred to as a “terminal” solution. The RA algorithm may also be modified to search for an integer feasible solution to a sub-problem using any of the known heuristics (e.g. Tabu Search, Simulated Annealing, Genetic Search), or to formulate and add Gomory’s cuts or other well know cuts (Wolsey, 1998) to sub-problems, change branching direction or priorities in the B&C algorithm. The optimization process outlined here doesn’t exclude these extensions.

Once a terminal solution is found, detailed tour schedules for agents can be developed in a number of ways. All such schedules will have the same objective value and, therefore, equally effective. First, work patterns and daily shift start times are specified. Consider a tour requiring consistent daily start times. The days-off scheduled in the solution for this tour are first assigned to individual tours. For example, if, in the terminal solution, tour type 1 has 2 agents from skill group 3 to start daily at 7:00am (that is, the variable $X^3_{1(7:00)} = 2$ in formulation (65-80)), initially these 2 agents are scheduled for work on all 7 days in a week. However, since day-off constraints (e.g. constraints (71-73)) ensures sufficient numbers of consecutive and non-consecutive days off for every agent, each of these two agents will get the required number of days-off. Suppose, for instance, each agent assigned to this tour receives only two consecutive days off. Then, in the solution, $\sum_h Y^3_{1(7:00)h} = 2$. Consequently, each agent will be off on one of

these two consecutive days off. Thus, the tour schedules for these two agents now have work and non-work days specified. Similarly, if a tour does not require consistent daily start time and shift length, daily shifts scheduled (including lengths and start times) in the terminal solution by the values of the decision variables are assigned to work patterns

5 specified by the values of the work pattern variables in the terminal solution. Work pattern constraints (e.g. constraint (77)) ensures that a daily shift is scheduled for every agent that will be at work due to the work patterns scheduled in the solution. These daily shifts are assigned to work patterns by following the variable definitions to satisfy tour and shift rules for each tour type. Thus, the weekly tours are stated with off days, work
10 days with daily start times and shift lengths for every agent scheduled in the solution. Break balance constraints (e.g. constraints (68-70) and (74-76)) ensure that one break of each type is scheduled for every agent scheduled to work on a given day. These breaks are assigned to respective tours that have a given day as a work day while making sure that agent group, tour type, the daily start time, shift length and break window
15 associations specified in the variable definitions are followed correctly to satisfy break scheduling requirements. Detailed weekly schedules are then assigned to specific agents based on their preferences, starting with the agent having the highest preference priority (e.g. seniority) to complete the rostering process.

20 Accordingly, it is to be understood that the embodiments of the invention herein described are merely illustrative of the application of the principles of the invention. It will be clear to one skilled in the art that the above embodiment may be altered or supplemented in many ways without departing from the scope of the invention. Reference herein to details of the illustrated embodiments are not intended to limit the
25 scope of claims, which themselves recite those features regarded as essential to the invention.

What is claimed is:

1. A computer implemented method for developing an optimized workforce schedule for a plurality of agents each with a combination of defined skills to serve one
30 or more contact types, a plurality of tour and shift scheduling rules located at

one or more contact centers each with its own operating hours and time zones,
comprising the steps of:

- (a) acquiring net staffing level requirements for each contact type, and for each period to be scheduled by any means;
- 5 (b) acquiring tour, shift, days-off, and break scheduling rules, agent skills groups, agent availability, and objective criterion to be optimized and its parameters by any means;
- (c) formulating the constraints and objective function of a Mixed Integer Programming (MILP) model to describe the tour and shift rules, staffing level requirements for a plurality of contact types and for each period to be
10 scheduled, agent skills, agent availability, and agent costs;
- (d) solving the LP relaxation of the MILP model formulated in the B&C algorithm, and stopping the B&C algorithm if an optimal solution to the MILP model is found;
- 15 (e) calling the RA algorithm comprising of the following steps if a solution to a node violating some integrality constraints is found by the B&C algorithm:
 - i. Obtaining the values of the decision variables in the solution found to the MILP model;
 - ii. Rounding the fractional values of decision variables U, W, V, QX, QU, QW, QV, and G down, and weekly tour variables X, and work pattern variables down if their fractional part is less than or equal to 0.50, and up if greater than 0.50, provided the agent availability constraints on the maximum number of agents available are not violated;
 - 20 iii. Scheduling additional days off by increasing the values of Y and Z if additional days off needed to satisfy tour scheduling requirements, and additional daily shifts to have a shift scheduled for every agent who will be working on a given day based of the work patterns scheduled;
 - 25 iv. Scheduling additional daily breaks if the number of breaks of each type is not sufficient to satisfy break scheduling rules for a tour, and
30 unscheduling breaks if more breaks are scheduled due to rounding;

- v. Computing agent shortages and excesses for each contact type and planning period;
 - 5 vi. Computing scheduled agent availability for all agent groups, comparing them with the agent allocations to different contact types by the rounded values of allocation variables G , and adjusting the values of the associated allocation variables to make agent allocations equal to scheduled agent availability;
 - 10 vii. Checking the solution constructed in steps (a) through (f) to find out if all agent requirements are met in every period and , when all requirements are met, eliminating all redundant agent tours and stopping with the integer feasible solution found;
 - viii. Stopping with a solution with agent shortages if agent requirements for some contact types are not met in all periods and all available agents who can serve these contact types are scheduled;
 - 15 ix. Continuing to step (i) if there are agents available to schedule and there shortages in some periods for some contact types;
 - x. Finding all periods in which some contact types have shortages, finding an agent with skills needed together with a complete tour schedule with work and non-work days, daily shift start times and shift lengths, daily break times to reduce the agent shortages, and adding to the solution by updating the values of the decision variables to include the newly found tour schedule;
 - 20 xi. Repeating steps (viii), (ix), and (x) until an integer feasible solution satisfying all requirements is found or the algorithm stops in step (viii);
 - 25 xii. Examining the tours scheduled in the integer feasible solution found and eliminating redundant tours
- (f) applying the RA algorithm to the solution found for the current node by the B&C algorithm, and passing it to the B&C algorithm if a solution better than
- 30 the best integer solution known by the RA algorithm is found;

2. The method of claim 1 further comprising of repeating steps (e) and (f) for every node the B&C algorithm solves and finds a solution violating only some integrality constraints, until a terminal solution to the MILP model is reached.

3. The method of claim 1 further comprising of processing the terminal solution found to assign daily shifts with start times and shift lengths to work patterns, and days-off scheduled to tours, and daily breaks to specific shifts to develop detailed weekly agent schedules;

4. The method of claim 1 further comprising of assigning agents to detailed weekly tours;

5. The method of claim 1, in which a terminal solution to the MILP model formulated is found when the objective function value for an integer feasible solution differs no more than a pre-specified percentage from the best lower bound found in the B&C algorithm for the MILP model.

6. The method of claim 1, in which a terminal solution to the MILP model formulated is found when an integer feasible solution is found and a pre-specified period of time is passed.

7. The method of claim 1, in which a terminal solution to the MILP model formulated is found when an integer feasible solution is found and a pre-specified number of nodes are solved in the B&C algorithm.

8. A computer program in a computer readable medium to be run on a computer for developing optimal schedules for a plurality of agents with a combination of defined skills, a plurality of contact types requiring defined agent skills, a plurality of tour and shift scheduling rules, and a plurality of contact centers, comprising of:

means for acquiring agent and skill requirements for each period to be scheduled;

means for acquiring, for each contact center to be scheduled, operating hours, agent skill groups, and scheduling paradigm for contact centers to specify whether each center is to be scheduled independently or a plurality of contact centers as a virtual contact center with combined;

means for acquiring tour and shift parameters and scheduling rules, and their availability to agent skills and work groups at different contact centers;

means for acquiring the objective criterion to be optimized and objective function

parameters to quantify cost or benefit;

means for generating an MILP model by defining decision variables, parameters and sets, and then formulating the objective function, agent and skill requirement constraints, breaks constraints, days-off and work pattern constraints, agent availability constraints, other schedule related constraints, and non-negativity and integrality constraints;

5

means for solving the MILP model using an optimization algorithm;

means for generating a detailed tour schedule using a terminal solution to the MILP model found by assigning daily shifts to work patterns scheduled,

10

days-off to weekly tours, and breaks to daily shifts;

means for assigning agents to specific tours.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:**

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.